

General

Title

Diabetes screening for people with schizophrenia or bipolar disorder who are using antipsychotic medications: percentage of members 18 to 64 years of age with schizophrenia or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

Source(s)

National Committee for Quality Assurance (NCQA). HEDIS 2016: Healthcare Effectiveness Data and Information Set. Vol. 1, narrative. Washington (DC): National Committee for Quality Assurance (NCQA); 2015. various p.

National Committee for Quality Assurance (NCQA). HEDIS 2016: Healthcare Effectiveness Data and Information Set. Vol. 2, technical specifications for health plans. Washington (DC): National Committee for Quality Assurance (NCQA); 2015. various p.

Measure Domain

Primary Measure Domain

Clinical Quality Measures: Process

Secondary Measure Domain

Does not apply to this measure

Brief Abstract

Description

This measure is used to assess the percentage of members 18 to 64 years of age with schizophrenia or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

Rationale

People with schizophrenia are at a greater risk of metabolic syndrome due to their serious mental illness

(Cohn et al., 2004). Diabetes screening is important for anyone with schizophrenia or bipolar disorder, and the added risk associated with antipsychotic medications contributes to the need to screen people with schizophrenia for diabetes. Diabetes screening for individuals with schizophrenia or bipolar disorder who are prescribed an antipsychotic medication may lead to earlier identification and treatment of diabetes.

Evidence for Rationale

Cohn T, Prud'homme D, Streiner D, Kameh H, Remington G. Characterizing coronary heart disease risk in chronic schizophrenia: high prevalence of the metabolic syndrome. *Can J Psychiatry*. 2004 Nov;49(11):753-60. [PubMed](#)

National Committee for Quality Assurance (NCQA). HEDIS 2015: Healthcare Effectiveness Data and Information Set. Vol. 1, narrative. Washington (DC): National Committee for Quality Assurance (NCQA); 2014. various p.

Primary Health Components

Schizophrenia; bipolar disorder; antipsychotic medication; diabetes screening; glucose test; hemoglobin A1c (HbA1c) test

Denominator Description

Medicaid members age 18 to 64 years as of December 31 of the measurement year with schizophrenia or bipolar disorder who were dispensed an antipsychotic medication (see the related "Denominator Inclusions/Exclusions" field)

Numerator Description

A glucose test or a hemoglobin A1c (HbA1c) test performed during the measurement year (see the related "Numerator Inclusions/Exclusions" field)

Evidence Supporting the Measure

Type of Evidence Supporting the Criterion of Quality for the Measure

A formal consensus procedure, involving experts in relevant clinical, methodological, public health and organizational sciences

One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Additional Information Supporting Need for the Measure

- In 2010, heart disease and diabetes were the leading causes of death in the United States (U.S.) (Murphy, Xu, & Kochanek, 2013). Because persons with serious mental illness who use antipsychotics are at increased risk of cardiovascular diseases and diabetes, screening and monitoring of these conditions is important.
- In 2007, diabetes was estimated to cost the U.S. economy \$174 billion. Of this, \$116 billion was attributed to medical care and \$58 billion to disability, work loss and premature death (Roger et al.,

2011).

- People with diabetes and schizophrenia or bipolar disorder have a 50 percent higher risk of death than diabetics without a mental illness (Vinogradova et al., 2010).
- Lack of appropriate care for diabetes and cardiovascular disease for people with schizophrenia or bipolar disorder who use antipsychotic medications can lead to worsening health and death. Addressing these physical health needs is an important way to improve health and economic outcomes downstream.

Evidence for Additional Information Supporting Need for the Measure

Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2010. Natl Vital Stat Rep. 2013 May 8;61(4):1-117.

National Committee for Quality Assurance (NCQA). The state of health care quality 2015. Washington (DC): National Committee for Quality Assurance (NCQA); 2015. 205 p.

Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, Carnethon MR, Dai S, de Simone G, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Greenlund KJ, Hailpern SM, Heit JA, Ho PM, Howard VJ, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM, Marelli A, Matchar DB, McDermott MM, Meigs JB, Moy CS, Mozaffarian D, Mussolino ME, Nichol G, Paynter NP, Rosamond WD, Sorlie PD, Stafford RS, Turan TN, Turner MB, Wong ND, Wylie-Rosett J. Heart disease and stroke statistics--2011 update: a report from the American Heart Association. Circulation. 2011 Feb 1;123(4):e18-209. [PubMed](#)

Vinogradova Y, Coupland C, Hippisley-Cox J, Whyte S, Penny C. Effects of severe mental illness on survival of people with diabetes. Br J Psychiatry. 2010 Oct;197(4):272-7. [PubMed](#)

Extent of Measure Testing

All HEDIS measures undergo systematic assessment of face validity with review by measurement advisory panels, expert panels, a formal public comment process and approval by the National Committee for Quality Assurance's (NCQA's) Committee on Performance Measurement and Board of Directors. Where applicable, measures also are assessed for construct validity using the Pearson correlation test. All measures undergo formal reliability testing of the performance measure score using beta-binomial statistical analysis.

Evidence for Extent of Measure Testing

Rehm B. (Assistant Vice President, Performance Measurement, National Committee for Quality Assurance, Washington, DC). Personal communication. 2015 Mar 16. 1 p.

State of Use of the Measure

State of Use

Current routine use

Current Use

not defined yet

Application of the Measure in its Current Use

Measurement Setting

Ambulatory/Office-based Care

Behavioral Health Care

Emergency Department

Hospital Inpatient

Hospital Outpatient

Managed Care Plans

Professionals Involved in Delivery of Health Services

not defined yet

Least Aggregated Level of Services Delivery Addressed

Single Health Care Delivery or Public Health Organizations

Statement of Acceptable Minimum Sample Size

Unspecified

Target Population Age

Age 18 to 64 years

Target Population Gender

Either male or female

National Strategy for Quality Improvement in Health Care

National Quality Strategy Aim

Better Care

National Quality Strategy Priority

Prevention and Treatment of Leading Causes of Mortality

Institute of Medicine (IOM) National Health Care Quality Report Categories

IOM Care Need

Living with Illness

IOM Domain

Effectiveness

Data Collection for the Measure

Case Finding Period

The measurement year

Denominator Sampling Frame

Enrollees or beneficiaries

Denominator (Index) Event or Characteristic

Clinical Condition

Encounter

Institutionalization

Patient/Individual (Consumer) Characteristic

Therapeutic Intervention

Denominator Time Window

not defined yet

Denominator Inclusions/Exclusions

Inclusions

Medicaid members age 18 to 64 years as of December 31 of the measurement year with schizophrenia or bipolar disorder who were dispensed an antipsychotic medication

Identify members with schizophrenia or bipolar disorder as those who met at least one of the following criteria during the measurement year:

At least one acute inpatient encounter, with any diagnosis of schizophrenia or bipolar disorder.

Any of the following code combinations meet criteria:

BH Stand Alone Acute Inpatient Value Set *with* Schizophrenia Value Set

BH Stand Alone Acute Inpatient Value Set *with* Bipolar Disorder Value Set

BH Stand Alone Acute Inpatient Value Set *with* Other Bipolar Disorder Value Set

BH Acute Inpatient Value Set *with* BH Acute Inpatient POS Value Set *and* Schizophrenia Value Set

BH Acute Inpatient Value Set *with* BH Acute Inpatient POS Value Set *and* Bipolar Disorder Value Set

BH Acute Inpatient Value Set *with* BH Acute Inpatient POS Value Set *and* Other Bipolar Disorder Value Set

At least two visits in an outpatient, intensive outpatient, partial hospitalization, emergency department (ED) or nonacute inpatient setting, on different dates of service, with any diagnosis of schizophrenia. Any two of the following code combinations meet criteria:

BH Stand Alone Outpatient/PH/IOP Value Set *with* Schizophrenia Value Set

BH Outpatient/PH/IOP Value Set *with* BH Outpatient/PH/IOP POS Value Set *and* Schizophrenia Value Set

ED Value Set *with* Schizophrenia Value Set

BH ED Value Set *with* BH ED POS Value Set *and* Schizophrenia Value Set

BH Stand Alone Nonacute Inpatient Value Set *with* Schizophrenia Value Set

BH Nonacute Inpatient Value Set *with* BH Nonacute Inpatient POS Value Set *and* Schizophrenia Value Set

At least two visits in an outpatient, intensive outpatient, partial hospitalization, ED or nonacute inpatient setting, on different dates of service, with any diagnosis of bipolar disorder. Any two of the following code combinations meet criteria:

BH Stand Alone Outpatient/PH/IOP Value Set *with* Bipolar Disorder Value Set

BH Stand Alone Outpatient/PH/IOP Value Set *with* Other Bipolar Disorder Value Set

BH Outpatient/PH/IOP Value Set *with* BH Outpatient/PH/IOP POS Value Set *and* Bipolar Disorder Value Set

BH Outpatient/PH/IOP Value Set *with* BH Outpatient/PH/IOP POS Value Set *and* Other Bipolar Disorder Value Set

ED Value Set *with* Bipolar Disorder Value Set

ED Value Set *with* Other Bipolar Disorder Value Set

BH ED Value Set *with* BH ED POS Value Set *and* Bipolar Disorder Value Set

BH ED Value Set *with* BH ED POS Value Set *and* Other Bipolar Disorder Value Set

BH Stand Alone Nonacute Inpatient Value Set *with* Bipolar Disorder Value Set

BH Stand Alone Nonacute Inpatient Value Set *with* Other Bipolar Disorder Value Set

BH Nonacute Inpatient Value Set *with* BH Nonacute Inpatient POS Value Set *and* Bipolar Disorder Value Set

BH Nonacute Inpatient Value Set *with* BH Nonacute Inpatient POS Value Set *and* Other Bipolar Disorder Value Set

Note:

Members must have been continuously enrolled during the measurement year.

Allowable Gap: No more than one gap in enrollment of up to 45 days during the measurement year. To determine continuous enrollment for a Medicaid beneficiary for whom enrollment is verified monthly, the member may not have more than a 1-month gap in coverage.

Exclusions

Exclude members who met any of the following criteria:

Members with diabetes. There are two ways to identify members with diabetes: by claim/encounter data and by pharmacy data. The organization must use both methods to identify members with diabetes, but a member need only be identified by one method to be excluded from the measure. Members may be identified as having diabetes during the measurement year or the year prior to the measurement year.

Claim/Encounter Data: Members who met any of the following criteria during the measurement year or the year prior to the measurement year (count services that occur over both years):

At least two outpatient visits (Outpatient Value Set), observation visits (Observation Value Set), ED visits (ED Value Set) or nonacute inpatient encounters (Nonacute Inpatient Value Set) on different dates of service, with a diagnosis of diabetes (Diabetes Value Set). Visit

type need not be the same for the two visits.

At least one acute inpatient encounter (Acute Inpatient Value Set) with a diagnosis of diabetes (Diabetes Value Set)

Pharmacy Data: Members who were dispensed insulin or oral hypoglycemics/antihyperglycemics during the measurement year or year prior to the measurement year on an ambulatory basis (refer to Table CDC-A in the original measure documentation for a list of prescriptions to identify members with diabetes).

Members who had no antipsychotic medications dispensed during the measurement year. There are two ways to identify dispensing events: by claim/encounter data and by pharmacy data. The organization must use both methods to identify dispensing events, but an event need only be identified by one method to be counted.

Claim/Encounter Data: An antipsychotic medication (Long-Acting Injections Value Set)

Pharmacy Data: Dispensed an antipsychotic medication (refer to Table SSD-D in the original measure documentation for a list of antipsychotic medications) on an ambulatory basis.

Value Set Information

Measure specifications reference value sets that must be used for HEDIS reporting. A value set is the complete set of codes used to identify the service(s) or condition(s) included in the measure. Refer to the [NCQA Web site](#) to purchase HEDIS Volume 2, which includes the Value Set Directory.

Exclusions/Exceptions

not defined yet

Numerator Inclusions/Exclusions

Inclusions

A glucose test (Glucose Tests Value Set) or a hemoglobin A1c (HbA1c) test (HbA1c Tests Value Set) during the measurement year, as identified by claim/encounter or automated laboratory data

Exclusions

Unspecified

Value Set Information

Measure specifications reference value sets that must be used for HEDIS reporting. A value set is the complete set of codes used to identify the service(s) or condition(s) included in the measure. Refer to the [NCQA Web site](#) to purchase HEDIS Volume 2, which includes the Value Set Directory.

Numerator Search Strategy

Fixed time period or point in time

Data Source

Administrative clinical data

Pharmacy data

Type of Health State

Does not apply to this measure

Instruments Used and/or Associated with the Measure

Unspecified

Computation of the Measure

Measure Specifies Disaggregation

Does not apply to this measure

Scoring

Rate/Proportion

Interpretation of Score

Desired value is a higher score

Allowance for Patient or Population Factors

not defined yet

Standard of Comparison

not defined yet

Identifying Information

Original Title

Diabetes screening for people with schizophrenia or bipolar disorder who are using antipsychotic medications (SSD).

Measure Collection Name

HEDIS 2016: Health Plan Collection

Measure Set Name

Effectiveness of Care

Measure Subset Name

Behavioral Health

Submitter

National Committee for Quality Assurance - Health Care Accreditation Organization

Developer

National Committee for Quality Assurance - Health Care Accreditation Organization

Funding Source(s)

Unspecified

Composition of the Group that Developed the Measure

National Committee for Quality Assurance's (NCQA's) Measurement Advisory Panels (MAPs) are composed of clinical and research experts with an understanding of quality performance measurement in the particular clinical content areas.

Financial Disclosures/Other Potential Conflicts of Interest

In order to fulfill National Committee for Quality Assurance's (NCQA's) mission and vision of improving health care quality through measurement, transparency and accountability, all participants in NCQA's expert panels are required to disclose potential conflicts of interest prior to their participation. The goal of this Conflict Policy is to ensure that decisions which impact development of NCQA's products and services are made as objectively as possible, without improper bias or influence.

Endorser

National Quality Forum - None

NQF Number

not defined yet

Date of Endorsement

2014 Dec 23

Adaptation

This measure was not adapted from another source.

Date of Most Current Version in NQMC

2015 Oct

Measure Maintenance

Unspecified

Date of Next Anticipated Revision

Unspecified

Measure Status

This is the current release of the measure.

This measure updates previous versions:

National Committee for Quality Assurance (NCQA). HEDIS 2015: Healthcare Effectiveness Data and Information Set. Vol. 1, narrative. Washington (DC): National Committee for Quality Assurance (NCQA); 2014. various p.

National Committee for Quality Assurance (NCQA). HEDIS 2015: Healthcare Effectiveness Data and Information Set. Vol. 2, technical specifications for health plans. Washington (DC): National Committee for Quality Assurance (NCQA); 2014. various p.

Measure Availability

Source available for purchase from the [National Committee for Quality Measurement \(NCQA\) Web site](#) .

For more information, contact NCQA at 1100 13th Street, NW, Suite 1000, Washington, DC 20005; Phone: 202-955-3500; Fax: 202-955-3599; Web site: www.ncqa.org .

Companion Documents

The following are available:

National Committee for Quality Assurance (NCQA). The state of health care quality 2015. Washington (DC): National Committee for Quality Assurance (NCQA); 2015 Oct. 205 p.

National Committee for Quality Assurance (NCQA). HEDIS 2016: Healthcare Effectiveness Data and Information Set. Vol. 2, technical update. Washington (DC): National Committee for Quality Assurance (NCQA); 2015 Oct 1. 12 p.

For more information, contact the National Committee for Quality Assurance (NCQA) at 1100 13th Street, NW, Suite 1000, Washington, DC 20005; Phone: 202-955-3500; Fax: 202-955-3599; Web site: www.ncqa.org .

NQMC Status

This NQMC summary was completed by ECRI Institute on July 25, 2013.

This NQMC summary was updated by ECRI Institute on January 22, 2014, April 10, 2015, and again on January 29, 2016.

Copyright Statement

This NQMC summary is based on the original measure, which is subject to the measure developer's copyright restrictions.

Content adapted and reproduced with permission from the National Committee for Quality Assurance (NCQA). HEDIS® is a registered trademark of NCQA. HEDIS measures and specifications were developed by and are owned and copyrighted by NCQA. HEDIS measures and specifications are not clinical guidelines

and do not establish a standard of medical care. NCQA makes no representations, warranties, or endorsement about the quality of any organization or physician that uses or reports performance measures and NCQA has no liability to anyone who relies on such measures or specifications. Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. NCQA disclaims all liability for use or accuracy of any coding contained in the specifications.

Anyone desiring to use or reproduce the measure abstracts without modification for a non-commercial purpose may do so without obtaining any approval from NCQA. All commercial uses of the measure abstracts must be approved by NCQA and are subject to a license at the discretion of NCQA. To purchase copies of the full measures and specifications, which contain additional distribution and use restrictions, contact NCQA Customer Support at 888-275-7585 or visit www.ncqa.org/publications

Production

Source(s)

National Committee for Quality Assurance (NCQA). HEDIS 2016: Healthcare Effectiveness Data and Information Set. Vol. 1, narrative. Washington (DC): National Committee for Quality Assurance (NCQA); 2015. various p.

National Committee for Quality Assurance (NCQA). HEDIS 2016: Healthcare Effectiveness Data and Information Set. Vol. 2, technical specifications for health plans. Washington (DC): National Committee for Quality Assurance (NCQA); 2015. various p.

Disclaimer

NQMC Disclaimer

The National Quality Measures Clearinghouse[®] (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the [NQMC Inclusion Criteria](#).

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. Moreover, the views and opinions of developers or authors of measures represented on this site do not necessarily state or reflect those of NQMC, AHRQ, or its contractor, ECRI Institute, and inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.